



*Promoting effective joint working by ME and CFS organisations to maximise impact on behalf of all people with ME and CFS in the UK*

We are representatives from Forward ME which is an advocacy group comprising charities, clinicians, and allied health professionals, led by the Countess of Mar and Carol Monaghan, MP.

Our common goal is to support people with Myalgic Encephalomyelitis (M.E.) and Post-viral fatigue syndrome (PVFS), and to raise awareness among other health care professionals and the wider community.

### **Post-Covid Syndrome**

A diagnosis of M.E. often follows a distinct and prolonged viral infection and a period of post-viral fatigue syndrome.

As you are no doubt aware there has been increasing discussion recently about a similar post-Covid syndrome where people still struggling with symptoms are also known as 'long-haulers.'

This is occurring in a significant minority of people who have survived the initial Covid-19 infection but who continue to experience debilitating symptoms; some of which appear to be unique to Covid-19, while others are remarkably like M.E.

Research has indicated that up to 10% of anyone who contracts a viral infection will experience a post-viral fatigue syndrome<sup>1</sup> and this figure may apply to covid-19 survivors as well.

As you are likely to encounter patients recovering from Covid-19 infection who will fall into this category, we wanted to alert you to the best practice approach to management and rehabilitation.

### **Post-Covid Rehabilitation**

Instead of the usual 'reconditioning', we recommend that these patients should be helped to adopt a paced approach to activity management.

This is used to avoid symptom exacerbation or post exertional malaise, a primary symptom of M.E., which can result in greater incapacity and even relapse. Similar issues also seem present in people struggling to recover from Covid infection.

The ME Association and The Royal College of Occupational Therapists have both published guidance on management and pacing for Post-Covid patients.<sup>2-4</sup>

Other strategies that might prove useful include ensuring individualised care and support, adequate rest, sleep, and nutrition. People suffering these lingering symptoms and the symptoms unique to Post-Covid will need appropriate help and ongoing care.

We believe it is possible to draw upon the experiences of patients with M.E. in effectively managing those with post-Covid syndrome. The following podcast features Dr Charles Shepherd from the ME Association and physiotherapists from 'Physios for ME'.

It discusses the rehabilitation needs of patients who are showing signs of post-viral and post-Covid syndrome and M.E.:

<https://www.youtube.com/watch?v=OyFNVayKYCg>



*Promoting effective joint working by ME and CFS organisations to maximise impact on behalf of all people with ME and CFS in the UK*

## **Graded Exercise Therapy**

It is important that patients receive the right care in a timely manner.

We feel it is vital that they do not receive graded exercise therapy (GET) as a matter of course because M.E. research and patient experience has shown it to be ineffective and harmful in many cases. This was highlighted in a survey commissioned by Forward ME and completed by an academic research group from Oxford Brookes University. It found that 80% of people with M.E. reported having adverse events because of GET<sup>5</sup>.

The report concluded, "GET is shown to cause considerable deterioration in physical and mental health". The National Institute of Health and Care Excellence (NICE) issued a statement cautioning against the use of GET in Post-Covid cases and noting that existing recommendations relating to M.E. were under review<sup>6</sup>.

The British Medical Journal (BMJ) also covered this announcement saying that, "Graded exercise therapy may not be appropriate for treating post-viral fatigue in patients recovering from covid-19"<sup>7</sup>.

## **Abnormal responses to exercise in ME**

There is now extensive evidence to show that people with M.E. have abnormal responses to exercise and while we don't yet know if this might apply to post-Covid survivors, it is possible there will be similarities and we would urge caution with any exercise as a precaution.

These abnormalities include reduced maximum heart rate<sup>8-10</sup>, reduced maximum oxygen consumption<sup>10-12</sup>, reduced cardiac output<sup>8,9,13</sup>, insufficient blood pressure increase on exertion<sup>11,14</sup>, decreased capacity to use oxygen<sup>8</sup>. anaerobic threshold and maximum exercise are reached at much lower oxygen capacity<sup>10,15</sup>, exhaustion reached more rapidly and accompanied by relatively reduced intracellular concentrations of ATP<sup>16</sup>, increased intracellular acidosis in exercising muscles and reduced post-exercise recovery from acidosis<sup>17,18</sup>.

Also, activation and worsening of symptoms which can be immediate or delayed by several days<sup>19,20</sup> – when exercise is repeated the next day, abnormalities are more severe<sup>21</sup> – decreased cognitive functioning and prolonged reaction time<sup>22</sup>, and a prolonged recovery period – usually 24 hours, often 48 but can last days, weeks or cause a relapse<sup>8,21,23</sup>.

We hope that the information provided is helpful and we would be grateful if you could disseminate this information to all clinicians and allied health professionals where appropriate.

Yours sincerely,

*On behalf of Forward ME*



Countess Margaret of Mar  
**Chair of Forward ME**



Carol Monaghan,  
**Deputy Chair of Forward ME**  
**Chair of the APPG on ME**  
**MP for Glasgow North-West**



Todd E Davenport PT, DPT,  
MPH, OCS  
**Professor & Program Director**  
**University of the Pacific**



Dr Nina Muirhead  
**NHS Dermatology Surgeon**  
**Chair of the CMRC Medical**  
**Education Group**

Dr William Weir  
**Infectious Disease Consultant**

Dr K.N. Hng  
**NHS Junior Doctor**



Bill and Janice Kent  
**ReMEMber**



Christine Harrison  
**Brame**



Tony Crouch  
**25% ME Group and Tymes Trust**

Natalie Hilliard  
Dr Nicola Clague-Baker  
Dr Michelle Bull  
Karen Leslie  
**Physios for ME**



Dr Charles Shepherd  
**Hon. Medical Adviser**  
Russell Fleming  
**The ME Association**



Dr Nigel Speight  
**Hon. Medical Adviser**  
**The ME Association and**  
**25% ME Group**



Sonya Chowdhury  
**Chief Executive Officer**  
Sam Bromiley  
Clare Ogden  
**Action for ME**



Jonathan Davies  
Sue Waddle  
**ME Research UK**



Dr Paul Worthley  
Helen Winning  
**Chief Executive**  
**The ME Trust**



Jane Colby  
**Executive Director**  
Debbie Burgess  
**Tymes Trust**



### Further reading and references:

1. Hickie I et al. Post-infective and chronic fatigue syndromes precipitated by viral and non-viral pathogens: prospective cohort study. *British Medical Journal*, 2006, 333 (7568): 575
2. ME Association free guidance on Post-Viral Syndrome and Post-Covid Syndrome: <https://meassociation.org.uk/wp-content/uploads/MEA-Covid-19-PVF-PVFS-MECFS-April-2020.pdf>
3. The Royal College of Occupational Therapists free guidance on Post-Covid Syndrome: <https://www.rcot.co.uk/recovering-covid-19-post-viral-fatigue-and-conserving-energy>
4. ME Association full range of free leaflets on Covid-19, Post-Covid, and ME: <https://meassociation.org.uk/product-category/covid-19/>



*Promoting effective joint working by ME and CFS organisations to maximise impact on behalf of all people with ME and CFS in the UK*

5. ME Research UK. Forward-ME survey on patients' experiences of CBT and GET; 2019 [cited 2019 Dec 11]. Available from: <https://www.meresearch.org.uk/cbt-and-get-survey/>
6. <https://www.nice.org.uk/guidance/gid-ng10091/documents/statement>
7. Torjesen I. British Medical Journal 2020;370:m2912
8. De Becker P, Roeykens J, Reynders M, et al. Exercise capacity in chronic fatigue syndrome. Arch Intern Med 2000;160:3270-77. [PMID: 11088089]
9. Inbar O, Dlin R, Rotstein A, Whipp BJ. Physiological responses to incremental exercise in patients with chronic fatigue syndrome. Med Sci Sports Exerc 2001; 33: 1463-70. [PMID: 11528333]
10. Jones DE, Hollingsworth KG, Jakovljevic DG, Fattakhova G, Pairman J, Blamire AM, Trenell MI, Newton JL. Loss of capacity to recover from acidosis on repeat exercise in chronic fatigue syndrome: a case-control study. Eur J Clin Invest 2012; 42: 186-94.. [PMID: 21749371]
11. Farquhar WB, Hunt BE, Taylor JA, Darling SE, Freeman R. Blood volume and its relation to peak O<sub>2</sub> consumption and physical activity in patients with chronic fatigue. Am J Physiol Heart Circ Physiol 2002; 282: H66-71. [PMID: 11748048]
12. Jammes Y, Steinberg JG, Mambrini O, Brégeon F, Delliaux S. Chronic fatigue syndrome: assessment of increased oxidative stress and altered muscle excitability in response to incremental exercise. J Intern Med 2005; 257: 299-310. [PMID: 15715687]
13. Peckerman A, La Manca JJ, Dahl KA, Chemitiganti R, Qureishi B, Natelson BH. Abnormal impedance cardiography predicts symptom severity in chronic fatigue syndrome. Am J Med Sci 2003; 326: 55-60. [PMID: 12920435]
14. Streeten DH. Role of impaired lower-limb venous innervation in the pathogenesis of the chronic fatigue syndrome. Am J Med Sci 2001;321:163-7.
15. Vermeulen RCW, Kurk RM, Visser FC, Sluiter W, Scholte HR. Patients with chronic fatigue syndrome performed worse than controls in a controlled repeated exercise study despite a normal oxidative phosphorylation capacity. J Transl Med 2010; 8: 93. [PMID: 20937116]
16. Wong R, Lopaschuk G, Zhu G, Walker D, Catellier D, Burton D, Teo K, Collins-Nakai R, Montague T. Skeletal muscle metabolism in the chronic fatigue syndrome. In vivo assessment by <sup>31</sup>P nuclear magnetic resonance spectroscopy. Chest. 1992; 102: 1716-22. [PMID: 1446478]
17. Chaudhuri A, Behan PO. In vivo magnetic resonance spectroscopy in chronic fatigue syndrome. Prostaglandins Leukot Essent Fatty Acids. 2004; 71: 181-3. [PMID: 15253888]
18. Jones DE, Hollingsworth KG, Taylor R, Blamire AM, Newton JL. Abnormalities in pH handling by peripheral muscle and potential regulation by the autonomic nervous system in chronic fatigue syndrome. J Intern Med 2010; 267: 394-401. [PMID: 20433583]
19. Yoshiuchi K, Farkas I, Natelson BH. Patients with chronic fatigue syndrome have reduced absolute cortical blood flow. Clin Physiol Funct Imaging 2006; 26: 83-6. [PMID: 16494597]



*Promoting effective joint working by ME and CFS organisations to maximise impact on behalf of all people with ME and CFS in the UK*

20. VanNess JM, Stevens SR, Bateman L, Stiles TL, Snell CR. Postexertional malaise in women with chronic fatigue syndrome. *J Womens Health (Larchmt)* 2010; 19: 239-244. [PMID: 20095909]
21. Van Oosterwijck J, Nijs J, Meeus M, Lefever I, Huybrechts L, et al. Pain inhibition and postexertional malaise in myalgic encephalomyelitis/chronic fatigue syndrome; an experimental study. *J Intern Med* 2010; 268: 265-78. [PMID: 20412374]
22. Manca JJ, Sisto SA, DeLuca J, Johnson SK, Lange G, Pareja J, Cook S, Natelson BH. Influence of exhaustive treadmill exercise on cognitive functioning in chronic fatigue syndrome. *Am J Med* 1998; 105: 59S-65S. [PMID: 9790484]
23. VanNess JM, Snell CR, Stevens SR. Diminished cardiopulmonary capacity during post-exertional malaise. *J Chronic Fatigue Syndr* 2007; 14: 77-85.